



© TESTOSTERONE UNDECANOATE CAPSULES

- Oral Testosterone Replacement approved by FDA (Testosterone Undecanoate)
- 96% Effective on Phase 3 Trial patients who completed on Day 90
- Most common side effect: ABPM SBP rise after 6 months (1.8 mmHG)
- Absorption through lymphatic pathway primarily bypassing liver. KYZATREX™ not known to cause adverse hepatic events.
- Absorption not significantly affected by food fat content or alcohol consumption
- Flexible Dosing: Easy to remember strengths (100, 150, 200mg)
- Titration Testing - Blood test flexibility of 3-5 hours after morning dose

Cash Pay: All Strengths
(No PA, No Adjudication, No Insurance Hassles)

Symptoms of Low Testosterone

Low Muscle Mass • Weight Gain • Fatigue/Low Energy • Low Sex Drive



Decreased Bone Density • Difficulty Sleeping • Depression/Anxiety



1. Submit eRx through EMR for fulfillment through:

DiRx Pharmacy

4754 N. Hiatus Rd, Sunrise, FL.
33351-7987

NPI: 1447555453

NCPDP: 5707801

2. Call in RX: (877) 367-3479

Fax in RX: (833) 347-9329

KYZATREX™ NDC Numbers

- NDC 80603-101-22 (100mg 120-count bottle)
- NDC 80603-103-22 (150mg 120-count bottle)
- NDC 80603-105-22 (200mg 120-count bottle)



For more information go to www.kyzatrex.com
or email care@dirxhealth.com



© TESTOSTERONE UNDECANOATE CAPSULES

IMPORTANT SAFETY INFORMATION

KYZATREX™ (testosterone undecanoate) is a prescription drug that is used to treat adult men who have low or no testosterone levels due to certain medical conditions.

KYZATREX™ is a controlled substance (CIII) because it contains testosterone. It is not known if KYZATREX™ is safe or effective in males younger than 18 years old. Improper use may affect bone growth in children. KYZATREX™ is not meant for use by women.

KYZATREX™ can increase blood pressure, which can increase the risk of having a heart attack or stroke and can increase risk of death due to a heart attack or stroke. Your risk may be greater if you have already had a heart attack or stroke or if you have other risk factors for heart attack or stroke.

- If your blood pressure increases while on KYZATREX™, blood pressure medicines may need to be started. If you are currently taking blood pressure medicines, they may need to be changed or new blood pressure medicines may need to be added to control your blood pressure.
- If your blood pressure cannot be controlled, KYZATREX™ may need to be stopped.
- Your healthcare provider will monitor your blood pressure while you are being treated with KYZATREX™.

Do not take KYZATREX™ if you:

- have breast cancer;
- have or might have prostate cancer;
- are a woman who is pregnant (KYZATREX™ may harm your unborn baby);
- are allergic to KYZATREX™ or any of its ingredients;
- or have low testosterone without certain medical conditions (e.g., do not take KYZATREX™ if you have low testosterone due to age).

Before you take KYZATREX™, tell your healthcare provider about all of your medical conditions, including if you:

- have high blood pressure or are treated for high blood pressure;
- have a history of diabetes;
- have heart problems;
- have high red blood cell count (hematocrit) or high hemoglobin laboratory value;
- have urinary problems due to an enlarged prostate;
- have liver or kidney problems;
- or have problems breathing while you sleep (sleep apnea).

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Taking KYZATREX™ with certain other medicines can affect each other. **Especially, tell your healthcare provider if you take:** insulin; medicines that decrease blood clotting (blood thinners); corticosteroids; or medicines that increase blood pressure, such as some cold medicine and pain medicines.

KYZATREX™ may cause other serious side effects including:

- **Increase in red blood cell count (hematocrit) or hemoglobin**, which can increase the risk of blood clots, strokes, and heart attacks. You may need to stop KYZATREX™ if your red blood cell count increases.
- **If you already have an enlarged prostate, your signs and symptoms may worsen while taking KYZATREX™.** These may include: increased urination at night; trouble starting your urine stream; urinating many times during the day; urge to go to the bathroom right away; a urine accident; inability to pass urine or weak urine flow.
- **Increased risk of prostate cancer.**
- **Blood clots in the legs or lungs.** Signs and symptoms of a blood clot in your leg can include pain, swelling or redness. Signs and symptoms of a blood clot in your lungs can include difficulty breathing or chest pain.
- **Abuse.** Testosterone can be abused when taken at higher than prescribed doses and when used with other anabolic androgenic steroids. Abuse can cause serious heart and psychological side effects.
- **In large doses, KYZATREX™ may lower your sperm count.**
- **Liver problems.** Symptoms of liver problems may include: nausea or vomiting; yellowing of your skin or whites of your eyes; dark urine; pain on the right side of your stomach area (abdominal pain).
- **Swelling of your ankles, feet, or body (edema), with or without heart failure.**
- **Enlarged or painful breasts.**
- **Breathing problems while you sleep (sleep apnea).**

Call your healthcare provider right away if you have any of the serious side effects listed above.

The most common side effect of KYZATREX™ is high blood pressure. Other side effects may include: headache, joint or back pain, diarrhea, increased red blood cell count, anxiety, constipation, swelling of the legs, and increased prostate specific antigen (PSA) levels.

These are not all the possible side effects of KYZATREX™. For more information, ask your healthcare provider or pharmacist.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088. You may also report side effects to Marius by visiting www.mariuspharma.com.

Keep KYZATREX™ and all medicines out of the reach of children.

See Full Prescribing Information and Medication Guide for KYZATREX™.